

3.1 EXERCISES

For the following exercises, use **Equation 3.1** to find the slope of the secant line between the values x_1 and x_2 for each function $y = f(x)$.

1. $f(x) = 4x + 7$; $x_1 = 2$, $x_2 = 5$
2. $f(x) = 8x - 3$; $x_1 = -1$, $x_2 = 3$
3. $f(x) = x^2 + 2x + 1$; $x_1 = 3$, $x_2 = 3.5$
4. $f(x) = -x^2 + x + 2$; $x_1 = 0.5$, $x_2 = 1.5$
5. $f(x) = \frac{4}{3x-1}$; $x_1 = 1$, $x_2 = 3$
6. $f(x) = \frac{x-7}{2x+1}$; $x_1 = 0$, $x_2 = 2$
7. $f(x) = \sqrt{x}$; $x_1 = 1$, $x_2 = 16$
8. $f(x) = \sqrt[3]{x-9}$; $x_1 = 10$, $x_2 = 13$
9. $f(x) = x^{1/3} + 1$; $x_1 = 0$, $x_2 = 8$
10. $f(x) = 6x^{2/3} + 2x^{1/3}$; $x_1 = 1$, $x_2 = 27$

For the following functions,

- a. use **Equation 3.4** to find the slope of the tangent line $m_{\tan} = f'(a)$, and
- b. find the equation of the tangent line to f at $x = a$.

11. $f(x) = 3 - 4x$, $a = 2$
12. $f(x) = \frac{x}{5} + 6$, $a = -1$
13. $f(x) = x^2 + x$, $a = 1$
14. $f(x) = 1 - x - x^2$, $a = 0$
15. $f(x) = \frac{7}{x}$, $a = 3$
16. $f(x) = \sqrt{x+8}$, $a = 1$
17. $f(x) = 2 - 3x^2$, $a = -2$
18. $f(x) = \frac{-3}{x-1}$, $a = 4$

19. $f(x) = \frac{2}{x+3}$, $a = -4$
20. $f(x) = \frac{3}{x^2}$, $a = 3$

For the following functions $y = f(x)$, find $f'(a)$ using **Equation 3.1**.

21. $f(x) = 5x + 4$, $a = -1$
22. $f(x) = -7x + 1$, $a = 3$
23. $f(x) = x^2 + 9x$, $a = 2$
24. $f(x) = 3x^2 - x + 2$, $a = 1$
25. $f(x) = \sqrt{x}$, $a = 4$
26. $f(x) = \sqrt{x-2}$, $a = 6$
27. $f(x) = \frac{1}{x}$, $a = 2$
28. $f(x) = \frac{1}{x-3}$, $a = -1$
29. $f(x) = \frac{1}{x^3}$, $a = 1$
30. $f(x) = \frac{1}{\sqrt{x}}$, $a = 4$

For the following exercises, given the function $y = f(x)$,

- a. find the slope of the secant line PQ for each point $Q(x, f(x))$ with x value given in the table.
- b. Use the answers from a. to estimate the value of the slope of the tangent line at P .
- c. Use the answer from b. to find the equation of the tangent line to f at point P .

31. [T] $f(x) = x^2 + 3x + 4$, $P(1, 8)$ (Round to 6 decimal places.)

x	Slope m_{PQ}	x	Slope m_{PQ}
1.1	(i)	0.9	(vii)
1.01	(ii)	0.99	(viii)
1.001	(iii)	0.999	(ix)
1.0001	(iv)	0.9999	(x)
1.00001	(v)	0.99999	(xi)
1.000001	(vi)	0.999999	(xii)

32. [T] $f(x) = \frac{x+1}{x^2-1}$, $P(0, -1)$

x	Slope m_{PQ}	x	Slope m_{PQ}
0.1	(i)	-0.1	(vii)
0.01	(ii)	-0.01	(viii)
0.001	(iii)	-0.001	(ix)
0.0001	(iv)	-0.0001	(x)
0.00001	(v)	-0.00001	(xi)
0.000001	(vi)	-0.000001	(xii)

33. [T] $f(x) = 10e^{0.5x}$, $P(0, 10)$ (Round to 4 decimal places.)

x	Slope m_{PQ}
-0.1	(i)
-0.01	(ii)
-0.001	(iii)
-0.0001	(iv)
-0.00001	(v)
-0.000001	(vi)

34. [T] $f(x) = \tan(x)$, $P(\pi, 0)$

x	Slope m_{PQ}
3.1	(i)
3.14	(ii)
3.141	(iii)
3.1415	(iv)
3.14159	(v)
3.141592	(vi)

[T] For the following position functions $y = s(t)$, an object is moving along a straight line, where t is in seconds and s is in meters. Find

- the simplified expression for the average velocity from $t = 2$ to $t = 2 + h$;
- the average velocity between $t = 2$ and $t = 2 + h$, where (i) $h = 0.1$, (ii) $h = 0.01$, (iii) $h = 0.001$, and (iv) $h = 0.0001$; and
- use the answer from a. to estimate the instantaneous

velocity at $t = 2$ second.


35. $s(t) = \frac{1}{3}t + 5$

36. $s(t) = t^2 - 2t$


37. $s(t) = 2t^3 + 3$

38. $s(t) = \frac{16}{t^2} - \frac{4}{t}$

39. Use the following graph to evaluate a. $f'(1)$ and b. $f'(6)$.



40. Use the following graph to evaluate a. $f'(-3)$ and b. $f'(1.5)$.



For the following exercises, use the limit definition of derivative to show that the derivative does not exist at $x = a$ for each of the given functions.

41. $f(x) = x^{1/3}$, $x = 0$

42. $f(x) = x^{2/3}$, $x = 0$

43. $f(x) = \begin{cases} 1, & x < 1 \\ x, & x \geq 1 \end{cases}$, $x = 1$

44. $f(x) = \frac{|x|}{x}$, $x = 0$


45. [T] The position in feet of a race car along a straight track after t seconds is modeled by the function $s(t) = 8t^2 - \frac{1}{16}t^3$.

- Find the average velocity of the vehicle over the following time intervals to four decimal places:
 - $[4, 4.1]$
 - $[4, 4.01]$
 - $[4, 4.001]$
 - $[4, 4.0001]$
- Use a. to draw a conclusion about the instantaneous velocity of the vehicle at $t = 4$ seconds.

46. [T] The distance in feet that a ball rolls down an incline is modeled by the function $s(t) = 14t^2$, where t is seconds after the ball begins rolling.

- Find the average velocity of the ball over the following time intervals:
 - $[5, 5.1]$
 - $[5, 5.01]$
 - $[5, 5.001]$
 - $[5, 5.0001]$
- Use the answers from a. to draw a conclusion about the instantaneous velocity of the ball at $t = 5$ seconds.

47. Two vehicles start out traveling side by side along a straight road. Their position functions, shown in the following graph, are given by $s = f(t)$ and $s = g(t)$, where s is measured in feet and t is measured in seconds.



- Which vehicle has traveled farther at $t = 2$ seconds?
- What is the approximate velocity of each vehicle at $t = 3$ seconds?
- Which vehicle is traveling faster at $t = 4$ seconds?
- What is true about the positions of the vehicles at $t = 4$ seconds?

48. [T] The total cost $C(x)$, in hundreds of dollars, to produce x jars of mayonnaise is given by $C(x) = 0.000003x^3 + 4x + 300$.

- a. Calculate the average cost per jar over the following intervals:
 - i. $[100, 100.1]$
 - ii. $[100, 100.01]$
 - iii. $[100, 100.001]$
 - iv. $[100, 100.0001]$
- b. Use the answers from a. to estimate the average cost to produce 100 jars of mayonnaise.

49. [T] For the function $f(x) = x^3 - 2x^2 - 11x + 12$, do the following.

- a. Use a graphing calculator to graph f in an appropriate viewing window.
- b. Use the ZOOM feature on the calculator to approximate the two values of $x = a$ for which $m_{\tan} = f'(a) = 0$.

50. [T] For the function $f(x) = \frac{x}{1+x^2}$, do the following.

- a. Use a graphing calculator to graph f in an appropriate viewing window.
- b. Use the ZOOM feature on the calculator to approximate the values of $x = a$ for which $m_{\tan} = f'(a) = 0$.

51. Suppose that $N(x)$ computes the number of gallons of gas used by a vehicle traveling x miles. Suppose the vehicle gets 30 mpg.

- a. Find a mathematical expression for $N(x)$.
- b. What is $N(100)$? Explain the physical meaning.
- c. What is $N'(100)$? Explain the physical meaning.

52. [T] For the function $f(x) = x^4 - 5x^2 + 4$, do the following.

- a. Use a graphing calculator to graph f in an appropriate viewing window.
- b. Use the nDeriv function, which numerically finds the derivative, on a graphing calculator to estimate $f'(-2)$, $f'(-0.5)$, $f'(1.7)$, and $f'(2.718)$.

53. [T] For the function $f(x) = \frac{x^2}{x^2 + 1}$, do the following.

- a. Use a graphing calculator to graph f in an appropriate viewing window.
- b. Use the nDeriv function on a graphing calculator to find $f'(-4)$, $f'(-2)$, $f'(2)$, and $f'(4)$.