

# 4 | APPLICATIONS OF DERIVATIVES



**Figure 4.1** As a rocket is being launched, at what rate should the angle of a video camera change to continue viewing the rocket? (credit: modification of work by Steve Jurvetson, Wikimedia Commons)

## Chapter Outline

- 4.1 Related Rates
- 4.2 Linear Approximations and Differentials
- 4.3 Maxima and Minima
- 4.4 The Mean Value Theorem
- 4.5 Derivatives and the Shape of a Graph
- 4.6 Limits at Infinity and Asymptotes
- 4.7 Applied Optimization Problems
- 4.8 L'Hôpital's Rule
- 4.9 Newton's Method
- 4.10 Antiderivatives

## Introduction

A rocket is being launched from the ground and cameras are recording the event. A video camera is located on the ground at a certain distance from the launch pad. At what rate should the angle of inclination (the angle the camera makes with the

ground) change to allow the camera to record the flight of the rocket as it heads upward? (See **Example 4.3**.)

A rocket launch involves two related quantities that change over time. Being able to solve this type of problem is just one application of derivatives introduced in this chapter. We also look at how derivatives are used to find maximum and minimum values of functions. As a result, we will be able to solve applied optimization problems, such as maximizing revenue and minimizing surface area. In addition, we examine how derivatives are used to evaluate complicated limits, to approximate roots of functions, and to provide accurate graphs of functions.

## 4.1 | Related Rates

### Learning Objectives

- 4.1.1** Express changing quantities in terms of derivatives.
- 4.1.2** Find relationships among the derivatives in a given problem.
- 4.1.3** Use the chain rule to find the rate of change of one quantity that depends on the rate of change of other quantities.

We have seen that for quantities that are changing over time, the rates at which these quantities change are given by derivatives. If two related quantities are changing over time, the rates at which the quantities change are related. For example, if a balloon is being filled with air, both the radius of the balloon and the volume of the balloon are increasing. In this section, we consider several problems in which two or more related quantities are changing and we study how to determine the relationship between the rates of change of these quantities.


### Setting up Related-Rates Problems

In many real-world applications, related quantities are changing with respect to time. For example, if we consider the balloon example again, we can say that the rate of change in the volume,  $V$ , is related to the rate of change in the radius,  $r$ . In this case, we say that  $\frac{dV}{dt}$  and  $\frac{dr}{dt}$  are **related rates** because  $V$  is related to  $r$ . Here we study several examples of related quantities that are changing with respect to time and we look at how to calculate one rate of change given another rate of change.

#### Example 4.1

##### Inflating a Balloon

A spherical balloon is being filled with air at the constant rate of  $2 \text{ cm}^3/\text{sec}$  (**Figure 4.2**). How fast is the radius increasing when the radius is  $3 \text{ cm}$ ?



**Figure 4.2** As the balloon is being filled with air, both the radius and the volume are increasing with respect to time.

##### Solution

The volume of a sphere of radius  $r$  centimeters is

$$V = \frac{4}{3}\pi r^3 \text{ cm}^3.$$

Since the balloon is being filled with air, both the volume and the radius are functions of time. Therefore,  $t$  seconds after beginning to fill the balloon with air, the volume of air in the balloon is

$$V(t) = \frac{4}{3}\pi[r(t)]^3 \text{ cm}^3.$$

Differentiating both sides of this equation with respect to time and applying the chain rule, we see that the rate of change in the volume is related to the rate of change in the radius by the equation

$$V'(t) = 4\pi[r(t)]^2 r'(t).$$

The balloon is being filled with air at the constant rate of  $2 \text{ cm}^3/\text{sec}$ , so  $V'(t) = 2 \text{ cm}^3/\text{sec}$ . Therefore,

$$2 \text{ cm}^3/\text{sec} = (4\pi[r(t)]^2 \text{ cm}^2) \cdot (r'(t) \text{ cm/s}),$$

which implies

$$r'(t) = \frac{1}{2\pi[r(t)]^2} \text{ cm/sec}.$$

When the radius  $r = 3 \text{ cm}$ ,

$$r'(t) = \frac{1}{18\pi} \text{ cm/sec}.$$



**4.1** What is the instantaneous rate of change of the radius when  $r = 6 \text{ cm}$ ?

Before looking at other examples, let's outline the problem-solving strategy we will be using to solve related-rates problems.

### Problem-Solving Strategy: Solving a Related-Rates Problem

1. Assign symbols to all variables involved in the problem. Draw a figure if applicable.
2. State, in terms of the variables, the information that is given and the rate to be determined.
3. Find an equation relating the variables introduced in step 1.
4. Using the chain rule, differentiate both sides of the equation found in step 3 with respect to the independent variable. This new equation will relate the derivatives.
5. Substitute all known values into the equation from step 4, then solve for the unknown rate of change.

Note that when solving a related-rates problem, it is crucial not to substitute known values too soon. For example, if the value for a changing quantity is substituted into an equation before both sides of the equation are differentiated, then that quantity will behave as a constant and its derivative will not appear in the new equation found in step 4. We examine this potential error in the following example.

## Examples of the Process

Let's now implement the strategy just described to solve several related-rates problems. The first example involves a plane flying overhead. The relationship we are studying is between the speed of the plane and the rate at which the distance between the plane and a person on the ground is changing.


## Example 4.2

### An Airplane Flying at a Constant Elevation

An airplane is flying overhead at a constant elevation of 4000 ft. A man is viewing the plane from a position 3000 ft from the base of a radio tower. The airplane is flying horizontally away from the man. If the plane is flying at the rate of 600 ft/sec, at what rate is the distance between the man and the plane increasing when the plane passes over the radio tower?

#### Solution

Step 1. Draw a picture, introducing variables to represent the different quantities involved.



**Figure 4.3** An airplane is flying at a constant height of 4000 ft. The distance between the person and the airplane and the person and the place on the ground directly below the airplane are changing. We denote those quantities with the variables  $s$  and  $x$ , respectively.

As shown,  $x$  denotes the distance between the man and the position on the ground directly below the airplane. The variable  $s$  denotes the distance between the man and the plane. Note that both  $x$  and  $s$  are functions of time. We do not introduce a variable for the height of the plane because it remains at a constant elevation of 4000 ft. Since an object's height above the ground is measured as the shortest distance between the object and the ground, the line segment of length 4000 ft is perpendicular to the line segment of length  $x$  feet, creating a right triangle.

Step 2. Since  $x$  denotes the horizontal distance between the man and the point on the ground below the plane,  $dx/dt$  represents the speed of the plane. We are told the speed of the plane is 600 ft/sec. Therefore,  $\frac{dx}{dt} = 600$  ft/sec. Since we are asked to find the rate of change in the distance between the man and the plane when the plane is directly above the radio tower, we need to find  $ds/dt$  when  $x = 3000$  ft.

Step 3. From the figure, we can use the Pythagorean theorem to write an equation relating  $x$  and  $s$ :

$$[x(t)]^2 + 4000^2 = [s(t)]^2.$$

Step 4. Differentiating this equation with respect to time and using the fact that the derivative of a constant is zero, we arrive at the equation

$$x \frac{dx}{dt} = s \frac{ds}{dt}.$$

Step 5. Find the rate at which the distance between the man and the plane is increasing when the plane is directly over the radio tower. That is, find  $\frac{ds}{dt}$  when  $x = 3000$  ft. Since the speed of the plane is 600 ft/sec, we know that  $\frac{dx}{dt} = 600$  ft/sec. We are not given an explicit value for  $s$ ; however, since we are trying to find  $\frac{ds}{dt}$  when  $x = 3000$  ft, we can use the Pythagorean theorem to determine the distance  $s$  when  $x = 3000$  and the height is 4000 ft. Solving the equation

$$3000^2 + 4000^2 = s^2$$

for  $s$ , we have  $s = 5000$  ft at the time of interest. Using these values, we conclude that  $\frac{ds}{dt}$  is a solution of the equation

$$(3000)(600) = (5000) \cdot \frac{ds}{dt}.$$

Therefore,

$$\frac{ds}{dt} = \frac{3000 \cdot 600}{5000} = 360 \text{ ft/sec}.$$

*Note:* When solving related-rates problems, it is important not to substitute values for the variables too soon. For example, in step 3, we related the variable quantities  $x(t)$  and  $s(t)$  by the equation

$$[x(t)]^2 + 4000^2 = [s(t)]^2.$$

Since the plane remains at a constant height, it is not necessary to introduce a variable for the height, and we are allowed to use the constant 4000 to denote that quantity. However, the other two quantities are changing. If we mistakenly substituted  $x(t) = 3000$  into the equation before differentiating, our equation would have been

$$3000^2 + 4000^2 = [s(t)]^2.$$

After differentiating, our equation would become

$$0 = s(t) \frac{ds}{dt}.$$

As a result, we would incorrectly conclude that  $\frac{ds}{dt} = 0$ .



**4.2** What is the speed of the plane if the distance between the person and the plane is increasing at the rate of 300 ft/sec?

We now return to the problem involving the rocket launch from the beginning of the chapter.

### Example 4.3

#### Chapter Opener: A Rocket Launch




**Figure 4.4** (credit: modification of work by Steve Jurvetson, Wikimedia Commons)

A rocket is launched so that it rises vertically. A camera is positioned 5000 ft from the launch pad. When the rocket is 1000 ft above the launch pad, its velocity is 600 ft/sec. Find the necessary rate of change of the camera's angle as a function of time so that it stays focused on the rocket.

### Solution

Step 1. Draw a picture introducing the variables.



**Figure 4.5** A camera is positioned 5000 ft from the launch pad of the rocket. The height of the rocket and the angle of the camera are changing with respect to time. We denote those quantities with the variables  $h$  and  $\theta$ , respectively.

Let  $h$  denote the height of the rocket above the launch pad and  $\theta$  be the angle between the camera lens and the

ground.

Step 2. We are trying to find the rate of change in the angle of the camera with respect to time when the rocket is 1000 ft off the ground. That is, we need to find  $\frac{d\theta}{dt}$  when  $h = 1000$  ft. At that time, we know the velocity of the rocket is  $\frac{dh}{dt} = 600$  ft/sec.

Step 3. Now we need to find an equation relating the two quantities that are changing with respect to time:  $h$  and  $\theta$ . How can we create such an equation? Using the fact that we have drawn a right triangle, it is natural to think about trigonometric functions. Recall that  $\tan \theta$  is the ratio of the length of the opposite side of the triangle to the length of the adjacent side. Thus, we have

$$\tan \theta = \frac{h}{5000}.$$


This gives us the equation

$$h = 5000 \tan \theta.$$

Step 4. Differentiating this equation with respect to time  $t$ , we obtain

$$\frac{dh}{dt} = 5000 \sec^2 \theta \frac{d\theta}{dt}.$$

Step 5. We want to find  $\frac{d\theta}{dt}$  when  $h = 1000$  ft. At this time, we know that  $\frac{dh}{dt} = 600$  ft/sec. We need to determine  $\sec^2 \theta$ . Recall that  $\sec \theta$  is the ratio of the length of the hypotenuse to the length of the adjacent side. We know the length of the adjacent side is 5000 ft. To determine the length of the hypotenuse, we use the Pythagorean theorem, where the length of one leg is 5000 ft, the length of the other leg is  $h = 1000$  ft, and the length of the hypotenuse is  $c$  feet as shown in the following figure.



We see that

$$1000^2 + 5000^2 = c^2$$

and we conclude that the hypotenuse is

$$c = 1000\sqrt{26} \text{ ft.}$$

Therefore, when  $h = 1000$ , we have

$$\sec^2 \theta = \left( \frac{1000\sqrt{26}}{5000} \right)^2 = \frac{26}{25}.$$

Recall from step 4 that the equation relating  $\frac{d\theta}{dt}$  to our known values is

$$\frac{dh}{dt} = 5000 \sec^2 \theta \frac{d\theta}{dt}.$$

When  $h = 1000$  ft, we know that  $\frac{dh}{dt} = 600$  ft/sec and  $\sec^2 \theta = \frac{26}{25}$ . Substituting these values into the



previous equation, we arrive at the equation

$$600 = 5000\left(\frac{26}{25}\right)\frac{d\theta}{dt}.$$

Therefore,  $\frac{d\theta}{dt} = \frac{3}{26}$  rad/sec.



**4.3** What rate of change is necessary for the elevation angle of the camera if the camera is placed on the ground at a distance of 4000 ft from the launch pad and the velocity of the rocket is 500 ft/sec when the rocket is 2000 ft off the ground?

In the next example, we consider water draining from a cone-shaped funnel. We compare the rate at which the level of water in the cone is decreasing with the rate at which the volume of water is decreasing.


### Example 4.4

#### Water Draining from a Funnel

Water is draining from the bottom of a cone-shaped funnel at the rate of  $0.03 \text{ ft}^3/\text{sec}$ . The height of the funnel is 2 ft and the radius at the top of the funnel is 1 ft. At what rate is the height of the water in the funnel changing when the height of the water is  $\frac{1}{2}$  ft?

#### Solution

Step 1: Draw a picture introducing the variables.



**Figure 4.6** Water is draining from a funnel of height 2 ft and radius 1 ft. The height of the water and the radius of water are changing over time. We denote these quantities with the variables  $h$  and  $r$ , respectively.



Let  $h$  denote the height of the water in the funnel,  $r$  denote the radius of the water at its surface, and  $V$  denote the volume of the water.

Step 2: We need to determine  $\frac{dh}{dt}$  when  $h = \frac{1}{2}$  ft. We know that  $\frac{dV}{dt} = -0.03$  ft/sec.

Step 3: The volume of water in the cone is

$$V = \frac{1}{3}\pi r^2 h.$$

From the figure, we see that we have similar triangles. Therefore, the ratio of the sides in the two triangles is the same. Therefore,  $\frac{r}{h} = \frac{1}{2}$  or  $r = \frac{h}{2}$ . Using this fact, the equation for volume can be simplified to

$$V = \frac{1}{3}\pi\left(\frac{h}{2}\right)^2 h = \frac{\pi}{12}h^3.$$

Step 4: Applying the chain rule while differentiating both sides of this equation with respect to time  $t$ , we obtain

$$\frac{dV}{dt} = \frac{\pi}{4}h^2 \frac{dh}{dt}.$$

Step 5: We want to find  $\frac{dh}{dt}$  when  $h = \frac{1}{2}$  ft. Since water is leaving at the rate of  $0.03$  ft<sup>3</sup>/sec, we know that

$\frac{dV}{dt} = -0.03$  ft<sup>3</sup>/sec. Therefore,

$$-0.03 = \frac{\pi}{4}\left(\frac{1}{2}\right)^2 \frac{dh}{dt},$$

which implies

$$-0.03 = \frac{\pi}{16} \frac{dh}{dt}.$$

It follows that

$$\frac{dh}{dt} = -\frac{0.48}{\pi} = -0.153 \text{ ft/sec.}$$



**4.4** At what rate is the height of the water changing when the height of the water is  $\frac{1}{4}$  ft?