Chapter 6 Applications of Integration 6.4 Arc Length of a Curve and Surface Area

Section Exercises

For the following exercises, find the length of the functions over the given interval.

165. y = 5x from x = 0 to x = 2Answer: $2\sqrt{26}$

166. $y = -\frac{1}{2}x + 25$ from x = 1 to x = 4Answer: $\frac{3\sqrt{5}}{2}$

167. x = 4y from y = -1 to y = 1Answer: $2\sqrt{17}$

168. Pick an arbitrary linear function x = g(y) over any interval of your choice (y_1, y_2) . Determine the length of the function and then prove the length is correct by using geometry.

Answer: Answers may vary

169. Find the surface area of the volume generated when the curve $y = \sqrt{x}$ revolves around the *x*-axis from (1, 1) to (4, 2), as seen here.



170. Find the surface area of the volume generated when the curve $y = x^2$ revolves around the y – axis from (1, 1) to (3,9).



For the following exercises, find the lengths of the functions of x over the given interval. If you cannot evaluate the integral exactly, use technology to approximate it.

171.
$$y = x^{3/2} \operatorname{from}(0, 0)$$
 to $(1, 1)$
Answer: $\frac{13\sqrt{13} - 8}{27}$
172. $y = x^{2/3} \operatorname{from}(1, 1)$ to $(8, 4)$
Answer: $\frac{1}{27} (80\sqrt{10} - 13\sqrt{13})$
173. $y = \frac{1}{3} (x^2 + 2)^{3/2} \operatorname{from} x = 0$ to $x = 1$
Answer: $\frac{4}{3}$
174. $y = \frac{1}{3} (x^2 - 2)^{3/2} \operatorname{from} x = 2$ to $x = 4$
Answer: $\frac{50}{3}$
175. **[T]** $y = e^x$ on $x = 0$ to $x = 1$
Answer: 2.0035
176. $y = \frac{x^3}{3} + \frac{1}{4x} \operatorname{from} x = 1$ to $x = 3$
Answer: $\frac{53}{6}$

177.
$$y = \frac{x^4}{4} + \frac{1}{8x^2}$$
 from $x = 1$ to $x = 2$
123

Answer: $\frac{125}{32}$

178.
$$y = \frac{2x^{3/2}}{3} - \frac{x^{1/2}}{2}$$
 from $x = 1$ to $x = 4$
Answer: $\frac{31}{6}$

179.
$$y = \frac{1}{27} (9x^2 + 6)^{3/2}$$
 from $x = 0$ to $x = 2$
Answer: 10

180. **[T]** $y = \sin x$ on x = 0 to $x = \pi$ Answer: 3.8202

For the following exercises, find the lengths of the functions of y over the given interval. If you cannot evaluate the integral exactly, use technology to approximate it.

181.
$$y = \frac{5-3x}{4}$$
 from $y = 0$ to $y = 4$
Answer: $\frac{20}{3}$
182. $x = \frac{1}{2}(e^{y} + e^{-y})$ from $y = -1$ to $y = 1$
Answer: $e - \frac{1}{e}$
183. $x = 5y^{3/2}$ from $y = 0$ to $y = 1$
Answer: $\frac{1}{675}(229\sqrt{229} - 8)$
184. **[T]** $x = y^{2}$ from $y = 0$ to $y = 1$
Answer: 1.4789
185. $x = \sqrt{y}$ from $y = 0$ to $y = 1$
Answer: $\frac{1}{8}(4\sqrt{5} + \ln(9 + 4\sqrt{5}))$

186.
$$x = \frac{2}{3}(y^2 + 1)^{3/2}$$
 from $y = 1$ to $y = 3$
Answer: $\frac{58}{3}$
187. **[T]** $x = \tan y$ from $y = 0$ to $y = \frac{3}{4}$
Answer: 1.201
188. **[T]** $x = \cos^2 y$ from $y = -\frac{\pi}{2}$ to $y = \frac{\pi}{2}$
Answer: 3.8202
189. **[T]** $x = 4^y$ from $y = 0$ to $y = 2$
Answer: 15.2341
190. **[T]** $x = \ln(y)$ on $y = \frac{1}{e}$ to $y = e$
Answer: 3.1962

For the following exercises, find the surface area of the volume generated when the following curves revolve around the *x*-axis. If you cannot evaluate the integral exactly, use your calculator to approximate it.

191.
$$y = \sqrt{x}$$
 from $x = 2$ to $x = 6$
Answer: $\frac{49\pi}{3}$
192. $y = x^3$ from $x = 0$ to $x = 1$
Answer: $\frac{\pi}{27} (10\sqrt{10} - 1)$
193. $y = 7x$ from $x = -1$ to $x = 1$
Answer: $70\pi\sqrt{2}$
194. **[T]** $y = \frac{1}{x^2}$ from $x = 1$ to $x = 3$
Answer: 5.5169
195. $y = \sqrt{4 - x^2}$ from $x = 0$ to $x = 2$
Answer: 8π

196. $y = \sqrt{4 - x^2}$ from x = -1 to x = 1Answer: 8π

197.
$$y = 5x$$
 from $x = 1$ to $x = 5$
Answer: $120\pi\sqrt{26}$

198. **[T]**
$$y = \tan x$$
 from $x = -\frac{\pi}{4}$ to $x = \frac{\pi}{4}$

Answer: 7.67816

For the following exercises, find the surface area of the volume generated when the following curves revolve around the *y*-axis. If you cannot evaluate the integral exactly, use your calculator to approximate it.

199.
$$y = x^{2}$$
 from $x = 0$ to $x = 2$
Answer: $\frac{\pi}{6}(17\sqrt{17}-1)$
200. $y = \frac{1}{2}x^{2} + \frac{1}{2}$ from $x = 0$ to $x = 1$
Answer: $\frac{2\pi}{3}(2\sqrt{2}-1)$
201. $y = x+1$ from $x = 0$ to $x = 3$
Answer: $9\sqrt{2\pi}$
202. **[T]** $y = \frac{1}{x}$ from $x = \frac{1}{2}$ to $x = 1$
Answer: 5.01642
203. $y = \sqrt[3]{x}$ from $x = 1$ to $x = 27$
Answer: $\frac{10\sqrt{10\pi}}{27}(73\sqrt{73}-1)$
204. **[T]** $y = 3x^{4}$ from $x = 0$ to $x = 1$
Answer: 15.8264
205. **[T]** $y = \frac{1}{\sqrt{x}}$ from $x = 1$ to $x = 3$
Answer: 25.645

206. **[T]** $y = \cos x$ from x = 0 to $x = \frac{\pi}{2}$

```
Answer: 10.0745
```

207. The base of a lamp is constructed by revolving a quarter circle $y = \sqrt{2x - x^2}$ around the y-axis from x = 1 to x = 2, as seen here. Create an integral for the surface area of this curve and compute it.



Answer: 2π

208. A light bulb is a sphere with radius 1/2 in. with the bottom sliced off to fit exactly onto a cylinder of radius 1/4 in. and length 1/3 in., as seen here. The sphere is cut off at the bottom to fit exactly onto the cylinder, so the radius of the cut is 1/4 in. Find the surface area (not including the top or bottom of the cylinder).



209. **[T]** A lampshade is constructed by rotating y = 1/x around the *x*-axis from y = 1 to y = 2, as seen here. Determine how much material you would need to construct this lampshade—that is, the surface area—accurate to four decimal places.



Answer: 10.5017

- 210. **[T]** An anchor drags behind a boat according to the function $y = 24e^{-x/2} 24$, where y represents the depth beneath the boat and x is the horizontal distance of the anchor from the back of the boat. If the anchor is 23 ft below the boat, how much rope do you have to pull to reach the anchor? Round your answer to three decimal places. Answer: 26.850 ft
- 211. **[T]** You are building a bridge that will span 10 ft. You intend to add decorative rope in the shape of $y = 5 |\sin((x\pi)/5)|$, where x is the distance in feet from one end of the

bridge. Find out how much rope you need to buy, rounded to the nearest foot. Answer: 23 ft

For the following exercises, find the exact arc length for the following problems over the given interval.

212.
$$y = \ln(\sin x)$$
 from $x = \pi/4$ to $x = (3\pi)/4$. (*Hint:* Recall trigonometric identities.)
Answer: $y = 2\ln\left(\cot\left(\frac{\pi}{8}\right)\right)$

- 213. Draw graphs of $y = x^2$, $y = x^6$, and $y = x^{10}$. For $y = x^n$, as *n* increases, formulate a prediction on the arc length from (0, 0) to (1, 1). Now, compute the lengths of these three functions and determine whether your prediction is correct. Answer: 2
- 214. Compare the lengths of the parabola $x = y^2$ and the line x = by from (0, 0) to (b^2, b) as *b* increases. What do you notice? Answer: Answers may vary
- 215. Solve for the length of $x = y^2$ from (0, 0) to (1, 1). Show that $x = (1/2) y^2$ from (0, 0) to (2, 2) is twice as long. Graph both functions and explain why this is so. Answer: Answers may vary
- 216. **[T]** Which is longer between (1, 1) and (2, 1/2): the hyperbola y = 1/x or the graph of x + 2y = 3?
- Answer: $y = \frac{1}{x}$
- 217. Explain why the surface area is infinite when y = 1/x is rotated around the x-axis for $1 \le x < \infty$, but the volume is finite.
- Answer: For more information, look up Gabriel's Horn.

This file is copyright 2016, Rice University. All Rights Reserved.